GARTERSNAKE IDENTIFICATION IN WISCONSIN

BUTLER'S GARTERSNAKE, COMMON GARTERSNAKE, PLAINS GARTERSNAKE


The ranges of three gartersnake species (Butler's gartersnake, common gartersnake, plains gartersnake) overlap significantly in southeastern Wisconsin. Gartersnakes need to be carefully examined to determine the species. Species are distinguished from one another by the location of their lateral (side) stripes. Lateral stripes involve a variety of combinations of scale rows. Scale rows are horizontal rows of scales found along the length of the snake and are numbered starting above the rectangular ventral scale (see close-up photos below). Scale rows are counted diagonally.

BUTLER'S GARTERSNAKE

The lateral stripe on the Butler's gartersnake (*Thamnophis butleri*) involves upper part of scale row 2, all of scale row 3, and often part of scale row 4. The lateral stripe is typically broad and golden yellow.

Wisconsin Range: Dodge, Fond du Lac, Jefferson, Milwaukee, Ozaukee, Sheboygan, Washington, Waukesha Counties






COMMON GARTERSNAKE

The lateral stripe on the common gartersnake (*Thamnophis sirtalis*) may involve the upper half of scale row 1 and always involves rows 2 and 3. The lateral stripe is of varying widths and shades of yellow.






PLAINS GARTERSNAKE


The lateral stripe on the plains gartersnake (*Thamnophis radix*) only involves scale row 3 and part or all of scale row 4. The lateral stripe is typically narrow and pale yellow.

Wisconsin Range: southern and west-central Wisconsin



Additional photos to aid in identifying species:





Butler's gartersnakes (left) typically have unmarked (or lightly marked) labial scales below the eye, while plains gartersnakes (right) typically have labial scales bordered in black.

Last updated: April 2, 2013



